arXiv:1304.0392v1 [cond-mat.str-el] 1 Apr 2013

Topological phase transitions in spin-S ladders

Natalia Chepiga,[1] Frédéric Michaud,[1] and Frédéric Mila[1]

[1] *Institute of Theoretical Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland*

(Dated: April 2, 2013)

We show that spin-S ladders undergo, for antiferromagnetic rung coupling, a series of 4S topological phase transitions when the ratio of the leg to rung coupling changes from strongly antiferromagnetic to strongly ferromagnetic. These phase transitions are characterized by a change of Berry phase on the rungs, and, with appropriate boundary conditions, by the appearance and disappearance of edge states without the bulk gap ever closing. The various phases can be distinguished by the number of valence bond singlets on the rungs, which decreases from 2S for strong rung coupling to 0 for dominant leg coupling. In particular, the isotropic spin-3/2 ladder with equal leg and rung coupling is explicitly shown to be in the valence bond solid phase with one singlet per bond.

PACS numbers: 75.10.Jm,75.10.Pq,75.40.Mg

Topological aspects of matter have become one of the dominant themes in solid state physics. The characterization of quantum phases by topological invariants has been an extremely fruitful concept in the Quantum Hall Effect[1, 2], and it lies at the root of more recent developments such as topological insulators[3, 4]. A related question concerns the nature of the phase transitions between quantum phases with different topological properties. Quantum phase transitions[5] that involve a symmetry breaking are often continuous, and, even if both phases are gapped, the gap closes at the transition. By contrast, transitions that do not involve any symmetry breaking are usually first-order, and it is in principle possible to go continuously from one phase to the other by taking another path, as in the liquid-gas transition by going around the critical point. In the case of transitions between phases that are only distinguished by topological properties, much less is known. It has been recently argued that, for topological phases to be really different, the system must possess some symmetry to protect them[6]. If this is the case, i.e. if phases are really distinct, there is however no general theory of what happens at the transition. In particular, under which circumstances should one expect edge states to appear and the bulk gap to close is not known. Simple examples of topological phase transitions are clearly called for to make progress in understanding their general properties.

In this Letter, we show that spin-S ladders with antiferromagnetic rung coupling undergo a cascade of topological phase transitions characterized by a topological invariant and by the appearance and disappearance of edge states in a parameter range where the bulk gap never closes, and where, accordingly, no quantum phase transition takes place. Spin ladders are defined by the Hamiltonian

$$H^{\text{Ladder}} = J_{\parallel}\sum_i\sum_{\alpha=1,2}\mathbf{S}_{i,\alpha}\cdot\mathbf{S}_{i+1,\alpha}+J_{\perp}\sum_i\mathbf{S}_{i,1}\cdot\mathbf{S}_{i,2} \quad (1)$$

where i is the rung index, α the leg one, and $\mathbf{S}_{i,\alpha}$ are spin S operators. Throughout we will use the parametrization $J_{\parallel} = J\cos\theta$ and $J_{\perp} = J\sin\theta$. Over the last two decades, spin ladders have been investigated in great detail[7]. It has been shown early on that, for antiferromagnetic rung coupling, half-integer spin ladders are gapped if the number of legs is even and gapless if it odd while integer spin ladders are always gapped, and for the two-leg ladder, a consensus has emerged, based on numerical investigations and field theory arguments, that there is no quantum phase transition between the weak and strong rung coupling regimes. This is to be contrasted with the case of dimerized spin chains, in which a series of quantum phase transitions take place upon increasing the dimerization, but despite specific investigations aimed at finding similar quantum phase transitions in the spin-1 two-leg ladder[8], the evidence has imposed itself that the spin gap remains open all the way from weak to strong rung coupling, definitely excluding the presence of a quantum phase transition. The only phase transition that has been detected in spin-1/2 ladders with antiferromagnetic leg coupling takes place when the sign of the rung coupling changes from antiferromagnetic to ferromagnetic. At this transition, the gap closes, and the two topologically distinct singlet phases can be distinguished by the type of string-order parameter (even or odd) that exhibits long-range order[9–11].

The starting point of the present study is the investigation of the Berry phase on the rungs of the ladders. The Berry phase can be defined for any Hamiltonian $H(\phi)$ which depends periodically on a parameter ϕ[12]. If $|GS(\phi)\rangle$ denotes a single-valued ground state of $H(\phi)$, the Berry connection is defined by $A(\phi) = \langle GS(\phi)|\partial_\phi|GS(\phi)\rangle$, and the Berry phase is the integration of the Berry connection over a loop: $i\gamma = \oint A(\phi)d\phi$. A few years ago, Hatsugai [13] has shown that one can detect the presence of a singlet on a given bond of an antiferromagnet by calculating the Berry phase associated to a twist of the transverse component of the spin-spin interaction on this bond: $S_i^+S_j^- + S_i^-S_j^+ \rightarrow e^{i\phi}S_i^+S_j^- + e^{-i\phi}S_i^-S_j^+$.



Figure 1: (Color online) Phase diagrams of the spin-S ladders for S=1/2, 1, 3/2 and 2 as a function of θ with the convention $J_\perp = J \sin\theta$ and $J_\parallel = J\cos\theta$. They consist of four types of phases: a ferromagnetic (FM) phase ($\pi < \theta < 3\pi/2$), a Haldane phase ($3\pi/2 < \theta < 2\pi$), and two types of singlet phases for antiferromagnetic rung coupling ($0 < \theta < \pi$). These singlet phases are characterized by a topological invariant, the Berry phase on a rung, which is equal to 0 in the light (yellow) phases and π in the dark (blue) phases. The integer number reported in each phase corresponds to the number of valence-bond singlets according to the valence-bond interpretation (see main text). It takes values between 0 to 2S, leading to a total of 4S phase transitions for antiferromagnetic rung coupling. For S=1/2, 1 and 3/2, the critical values of θ are the result of a finite-size analysis, while for S=2 they correspond to the values for 8 sites.

In a valence bond solid, the Berry phase is related to the number N_s of singlets of the bond by:

$$\gamma = N_s\,\pi \pmod{2\pi}, \tag{2}$$

this quantization being protected by time-reversal symmetry.

This criterion has been tested on dimerized spin chains with larger spins[14], which undergo a series of quantum phase transitions upon increasing the dimerization[15], and indeed the Berry phase jumps between 0 and π at each transition, in agreement with the interpretation of the phases in terms of valence-bond singlets.

For spin ladders, since there is no quantum phase transition upon increasing the rung coupling, one might naively expect that the Berry phase does not change upon increasing the rung coupling, and that it is always equal to the value it has to take in the limit of strong rung coupling, π for half-integer spins with an odd number of valence-bond singlets and 0 for integer spins with an even number of valence-bond singlets. To investigate this point further, we have calculated the Berry phase for spin-S ladders with S=1/2, 1, 3/2 and 2 as a function of the rung coupling (see Supplemental Material for details). The calculations have been performed on finite ladders with periodic boundary conditions (PBC). It turns out that, for antiferromagnetic rung coupling, the Berry phase changes several times between 0 and π upon increasing the rung coupling for both antiferromagnetic and ferromagnetic leg coupling, resulting in the phase diagrams of Fig. 1. The critical values of θ have been deduced from a finite scaling analysis, except for S=2, where results for 8 sites are shown. The scaling is quite smooth, up to an even-odd effect for antiferromagnetic leg coupling, as can be seen from the example of the spin-1/2 ladder (see Fig. 2). The finite-size scalings for spin-1 and spin-3/2 ladders are shown in the Supplemental Material. Note that the transition at $\theta = 0.26\,\pi$ for spin 1/2 has already been reported in Ref. 16 in the context of an investigation of the frustrated spin-1/2 ladder with four-spin ring exchange. Note also that this transition does not show up in the string order parameters that distinguish the phases with ferromagnetic and antiferromagnetic rung couplings[9–11].

To be complete, we have also included the ferromagnetic rung coupling part of the phase diagram, which simply consists of two phases: a ferromagnetic phase when the leg coupling is ferromagnetic (see Supplemental Material), and a Haldane phase [17] when the leg coupling is antiferromagnetic. For strong ferromagnetic rung coupling, the spins of a rung bind into a spin 2S, and the low-energy effective model is just a spin-2S Heisenberg chain. This Haldane chain can be characterized by the Berry phase on one of its bonds, which is equal to π for 2S odd and 0 for 2S even. For the ladder, this requires to calculate the Berry phase associated to simultaneous twists on two leg bonds facing each other. We have found that this Berry phase is constant throughout that phase, with a value equal to π for half-integer spin (2S odd) and to 0 for integer spin (2S even), in agreement with the interpretation of this phase as a Haldane phase.

For $-\pi/2 < \theta < \pi$, the spin gap never closes for integer spins, and it only closes at the transition between the Haldane phase and the other singlet phases for half-integer spins (for $\theta = 0$, the system consists of decoupled half-integer spin chains, which are gapless). So the phase transitions detected by the Berry phase for $0 < \theta < \pi$ cannot be quantum phase transitions. Since the Berry phase is a topological invariant, they must be topological in nature. The next question is thus to determine the nature of the various phases. In Ref. 16 , the authors proposed the name "rung * singlet" for the weak-rung phase below $\theta = 0.26\,\pi$, suggesting that it is another type of rung singlet phase to be further investigated. However, in view of the cascade of phase transitions for



Figure 2: (Color online) Finite-size scaling of the topological transitions of the spin-1/2 ladder. The results have been fitted with a fourth order polynomial in $1/N$.

larger spins, it seems more natural, by analogy with the dimerized chains, to interpret the various phases as corresponding to different numbers of valence-bond singlets on the rungs, going from 2S in the strong rung limit, when pairs of spins form singlets on each rung, to 0 upon approaching the limit of decoupled chains. According to this interpretation, the weak-rung phase of the spin-1/2 ladder is not a rung singlet phase since it has no singlet on the rung.

To check this interpretation in terms of valence-bond singlets, we have investigated another possible characteristics of topological phase transitions, namely the appearance or disappearance of edge states at the transition. In the Haldane phase, it has long been known that edge states appear in finite chains[18–21]. In the present case, these edge states are expected to appear if we consider finite ladders with vertical edges, and indeed we have obtained numerical evidence of the presence of such edge states in the Haldane phase of the spin-1/2 ladder (not shown). By contrast, to distinguish the singlet phases for antiferromagnetic rung coupling, one should rather consider open boundary conditions (OBC) with diagonal edges[22]. Indeed, in the strong rung limit, the spins on the rungs are bound into singlets, and the orphan spin at each end of the ladder is expected to give rise to edge states. So we have investigated the fate of the edge states with diagonal edges as a function of the rung coupling in a spin-1/2 ladder. The low energy states with PBC and with diagonal edges are plotted as the inverse number of spins for two values of theta: $\theta = 0.3\,\pi$, where the Berry phase is equal to 1, and $\theta = 0.1\,\pi$, where the Berry phase is equal to 0. For PBC, the gap is finite in all cases, in agreement with previous results. For OBC with diagonal edges, edge states are indeed present for $\theta = 0.3\,\pi$, but they disappear when the Berry phase changes from π to 0, and they are not present any more for $\theta = 0.1\,\pi$. This confirms the interpretation of the change of Berry phase as a topological phase transition, with the disappearance of edge states upon reducing the rung coupling.

What is remarkable as compared to other cases where edge states appear is that there is neither a first-order transition (unlike the case of frustrated $S \geq 1$ spin chains[23, 24]), nor a continuous phase transition with a gap closing, a standard argument to explain the appearance of edge states at a topological phase transition relying on the gap closing at the transition upon going from one phase to the other. So, to better understand the fate of the edge states at the transition, we have calculated the evolution of the magnetization profile in the $S_z = 1$ sector of a ladder with diagonal boundary conditions from the strong rung limit to the transition at $\theta = 0.26\,\pi$. In the strong rung limit, the magnetization is concentrated at the edges of the system, and the fast decay can be fitted with an exponential with a very small length scale ξ^M (see Supplemental Material). Upon approaching the transition, the magnetization is less and less localized, a behavior consistent with a divergence of ξ^M. A similar information can be extracted from the energy spectrum: the energy of the edge states collapse onto the ground state exponentially with the inverse number of rungs, and the length ξ^E that can be extracted from the fit is in good agreement with ξ^M.

So, at the transition, a typical length scale seems to diverge. This is usually associated to a gap closing, but not here. This can be interpreted in the following way: At the transition, the spins that give rise to the edge states are no longer localized at the end of the chains. If they were free to move along the chain, they would give rise to a band of low-lying excitations and to a gapless spectrum. However, they move on a ladder, not on a chain, and this peculiar geometry opens a gap in their spectrum.

For larger spins, a complete picture of the edge states cannot be obtained on the basis of exact diagonalizations because of the size limitation, and it will be necessary to use more sophisticated numerical approaches, which goes beyond the scope of the present paper. Still, as shown in the Supplemental Material, clear indications of changes in the edge states at the transitions could be obtained for spin-1, where, in the strong-rung limit, we expect to have two orphan spins 1 at each edge, hence 9 low-lying states below the gap (a singlet, a triplet and a quintuplet).

As an alternative check of the valence-bond singlet picture, we have looked at the isotropic spin-3/2 ladder ($J_\perp = J_\parallel$). $\theta = \pi/4$ lies in a phase with Berry phase π, and the valence-bond singlet picture puts it in the phase with one singlet per rung. Since a spin 3/2 is able to build three valence-bond singlets, and since the coordinance of a 2-leg ladder is equal to 3, this phase is consistent with the valence-bond solid phase in which each bond (rungs and legs) carries one valence-bond singlet, i.e. in which the Berry phase of all bonds (and not just on the rungs) is equal to π. So we have calculated the Berry phase on the leg bonds, and indeed it is equal to π in that phase. So, it seems that the isotropic spin-3/2 ladder is a valence-bond solid.

To further confirm this conclusion, we have studied the



Figure 3: (Color online) Top panels: Low energy spectrum of the spin-$S = 1/2$ ladder with two types of boundary conditions (PBC and OBC with diagonal edges) for two values of the rung coupling: (a) in the rung phase ($\theta = 0.3\,\pi$), where an edge state is clearly present for OBC with diagonal edges and (b) in phase without valence-bond singlets on the rungs ($\theta = 0.1\pi$) where there is no edge state. Panel (c): Local magnetization in the sector $S^z_{\mathrm{tot}} = 1$ for the spin $S = 1/2$ ladder with diagonal edges. Inset: sketch of a cluster with diagonal edges. Squares (circles) have been used for sites of the top (bottom) leg. (d): Correlation length as a function of θ as deduced from the exponential scaling of the edge state gap (ξ^E) with the number of rungs $N_r \equiv N/2$, and from fitting the magnetization profile (ξ^M).

Hamiltonian $H(\alpha) = (1-\alpha)H^{\mathrm{Ladder}} + \alpha H^{\mathrm{AKLT}}$ that interpolates between the Heisenberg model and the AKLT model[25]:

$$H^{\mathrm{AKLT}} = J_{\parallel}\sum_i \sum_{\alpha=1,2} P_{(i,\alpha),(i+1,\alpha)} + J_{\perp}\sum_i P_{(i,1),(i,2)},$$

where $P_{i,j} = \frac{11}{128} + \frac{27}{160}\mathbf{S}_i\cdot\mathbf{S}_j + \frac{29}{360}(\mathbf{S}_i\cdot\mathbf{S}_j)^2 + \frac{1}{90}(\mathbf{S}_i\cdot\mathbf{S}_j)^3$ is the projector on total spin 3. When $\alpha = 1$, the AKLT state is the ground state for any θ in the range $0 < \theta < \pi/2$ since $J_{\parallel}$ and $J_{\perp}$ are strictly positive. As can be seen in Fig. 4, this AKLT phase at $\alpha = 1$ smoothly evolves into the intermediate phase with a rung Berry phase π and one valence-bond singlet per rung ($0.203\,\pi < \theta < 0.308\,\pi$) at $\alpha = 0$. The leg Berry phase (not shown) is equal to π in all this phase. This unambiguously demonstrates that the S=3/2 isotropic ladder is in the same phase as the AKLT model, hence that it is in the valence-bond solid phase.

In the same spirit, the valence-bond singlet picture predicts that the phase of the spin-2 ladder with two valence-bond singlets per rung should have one valence-bond singlet per leg bond, hence a Berry phase π, a prediction confirmed by our numerical results.



Figure 4: (Color online) Rung Berry phase of the model that interpolates between the Heisenberg and AKLT models as a function of α for 8 sites. The AKLT valence-bond solid at $\alpha = 1$ (sketched as an inset) evolves smoothly into the intermediate phase that contains the isotropic ladder at $\alpha = 0$.

To summarize, we have shown that spin-S ladders with antiferromagnetic rungs undergo a cascade of topological phase transitions upon changing the ratio of leg to rung coupling. At the transitions, a topological invariant, the Berry phase on a rung, changes from 0 to π, and edge states appear or disappear, but the bulk gap does not close. The absence of gap closing might be taken as an indication that there is no true phase transition. However, as first pointed out by Hatsugai[13], the quantization of the Berry phase as equal to 0 or π module 2π is a consequence of time reversal symmetry, and phases with Berry phase 0 and π are topologically distinct, the difference being protected by this symmetry in the spirit of Ref. 6. In that respect, the phases with the same Berry phase modulo $2\,\pi$, i.e. with the same parity of the number of singlets in the valence-bond singlet interpretation, should not be considered as topological distinct phases, although the classification according to the number of valence-bond singlets (and not simply its parity) is useful in discussing edge states.

So the present results lead to the conclusion that, at the transition between different topological phases, the bulk gap does not necessarily close, even if edge states appear. What happens is that edge states indeed delocalize at the transition, but the spectrum remains gapped because of another interaction, here the interaction between the chains due to the ladder geometry. It will be interesting to see if similar topological phase transitions without gap closing appear in other contexts.

We are indebted to E. Orignac and K. Penc for enlightening comments on the fate of the edge states at the transition. This work has been supported by the Swiss National Fund and by MaNEP.

[1] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Phys. Rev. Lett. **49**, 405 (1982).

[2] J. E. Avron, D. Osadchy, and R. Seiler, Physics Today **56**, 38 (2003).

[3] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).

[4] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).

[5] S. Sachdev, *Quantum Phase Transitions* (University Press, ADDRESS, 2001).

[6] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Phys. Rev. B **85**, 075125 (2012).

[7] E. Dagotto and T. Rice, Science **271**, 618 (1996).

[8] S. Todo, M. Matsumoto, C. Yasuda, and H. Takayama, Phys. Rev. B **64**, 224412 (2001).

[9] Y. Nishiyama, N. Hatano, and M. Suzuki, Journal of the Physical Society of Japan **64**, 1967 (1995).

[10] E. H. Kim, G. Fáth, J. Sólyom, and D. J. Scalapino, Phys. Rev. B **62**, 14965 (2000).

[11] G. Fáth, O. Legeza, and J. Sólyom, Phys. Rev. B **63**, 134403 (2001).

[12] M. V. Berry, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences **392**, 45 (1984).

[13] Y. Hatsugai, Journal of the Physical Society of Japan **75**, 123601 (2006).

[14] T. Hirano, H. Katsura, and Y. Hatsugai, Phys. Rev. B **77**, 094431 (2008).

[15] I. Affleck and F. D. M. Haldane, Phys. Rev. B **36**, 5291 (1987).

[16] I. Maruyama, T. Hirano, and Y. Hatsugai, Phys. Rev. B **79**, 115107 (2009).

[17] F. D. M. Haldane, Physics Letters A **93**, 464 (1983).

[18] T. Kennedy, Journal of Physics: Condensed Matter **2**, 5737 (1990).

[19] M. Hagiwara *et al.*, Phys. Rev. Lett. **65**, 3181 (1990).

[20] T.-K. Ng, Phys. Rev. B **50**, 555 (1994).

[21] P. Lecheminant and E. Orignac, Phys. Rev. B **65**, 174406 (2002).

[22] M. Arikawa, S. Tanaya, I. Maruyama, and Y. Hatsugai, Phys. Rev. B **79**, 205107 (2009).

[23] A. Kolezhuk, R. Roth, and U. Schollwöck, Phys. Rev. Lett. **77**, 5142 (1996).

[24] R. Roth and U. Schollwöck, Phys. Rev. B **58**, 9264 (1998).

[25] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).

SUPPLEMENTAL MATERIAL FOR “TOPOLOGICAL PHASE TRANSITIONS IN SPIN-S LADDERS”

NUMERICAL CALCULATION OF THE BERRY PHASE

In the main text, we have given a definition of the Berry phase as an integral of the Berry connection over a closed path:

$$i\gamma = \oint A(\phi)d\phi, \tag{3}$$

where $A(\phi) = \langle GS(\phi)|\partial_\phi|GS(\phi)\rangle$. To calculate the Berry phase numerically, it is expedient, following Ref. 1, to introduce a gauge-invariant Berry phase on the lattice by discretizing in ϕ:

$$\gamma_K = \sum_{k=0}^{K-1} \arg\langle GS(\phi_k)|GS(\phi_{k+1})\rangle, \quad \phi_k = \frac{2\pi k}{K} \tag{4}$$

The Berry phase is then given by $\gamma = \lim_{K\to\infty}\gamma_K$. In practice, the convergence is achieved very fast. Throughout, we have used $K = 16$.

Now, for large system sizes, it becomes numerically quite expensive to compute the Berry phase because one needs to diagonalize the system an important number of times (here 16), with just a change on one bond. However, a change of Berry phase is associated to a level crossing, and this level crossing usually appears for simple values of the twist parameter, $\phi = 0$ or $\phi = \pi$. Since we know that the gap never closes for $\phi = 0$, the level crossing has to happen for $\phi = \pi$, i.e. for anti-periodic boundary conditions. In other words, it is sufficient to compute the ground state energy of the system with anti-periodic boundary conditions, hence for a single value of ϕ, and to determine the parameter at which a level crossing occurs. This analysis is done in Fig. 5 for $N = 30$ sites around the first transition at $\theta \approx 0.26\,\pi$. The systematic investigation of large system sizes has been performed with this method, and not by computing explicitly the Berry phase.

FINITE-SIZE ANALYSIS OF THE TRANSITION POINTS FOR SPIN $S = 1$ AND $S = 3/2$ LADDERS

In Fig. 6, we show the finite-size analysis of the transitions for spin $S = 1$ and $S = 3/2$ ladders. The results for $S = 1/2$ can be found in the main text. The extrapolated values have been obtained by fitting the results with first or second order polynomials in $1/N$.

For spin $S = 1$, we did not observe any change of Berry phase for $N = 6$ and $N = 10$ for the first two transitions. This is an extreme case of the even-odd effect, which is generally present when the leg coupling is antiferromagnetic. So the fit has been performed using only the results for systems with an even number of rungs, i.e. $N = 8$, $N = 12$ and $N = 16$.



Figure 5: (Color online) Closing of the gap at the transition around $\theta \approx 0.26\,\pi$ for the spin-1/2 ladder with $N = 30$ sites and twisted periodic conditions. A very precise estimate of the transition point can be simply obtained by a linear fit.

For $S = 3/2$, the same is true for the transitions around $\theta \approx 0.308\,\pi$ and $\theta \approx 0.389\,\pi$. For $\theta \approx 0.203\,\pi$, we clearly see a strong even-odd effect. As in the case of the spin-1/2 ladder, extrapolating the results separately for systems with an even and odd number of rungs respectively leads to results in excellent agreement, with a difference of the order of 0.00005 in the present case.

EDGE STATES FOR THE SPIN $S = 1$ LADDER.

In this section, we summarize the results we have obtained regarding the appearance of edge states for spin-1 ladders under open boundary conditions with diagonal edges.

In the strong antiferromagnetic rung coupling limit, two isolated spins $S = 1$ appear at the edges, forming 9 low lying energy levels, one quintuplet, one triplet and one singlet. Since we work in the subspace $S_z^{\mathrm{tot}} = 0$, we expect to see only three states, namely the singlet, one of the triplets, and one of the quintuplets. The first one is the ground state. So, since we measure the energies with respect to the ground state energy, we expect to find two low-lying states. This is clearly the case in Fig. 7 (a), which corresponds to $\theta = 0.45\,\pi$, well inside the strong rung singlet phase with two valence-bond singlets per rung.

The situation is quite different in the intermediate phase ($\theta = 0.3\,\pi$) shown Fig. 7 (b). Indeed, open bound-



Figure 6: (Color online) Finite-size analysis of the topological transitions in the spin-1 (left panel) and spin-3/2 (right panel) ladders. The fits have been performed with first and second order polynomials in $1/N$

ary conditions with diagonal edges give rise to only one state below the gap. In addition, its energy is not so small. With the system sizes available, it is not possible to decide whether its energy goes to zero in the thermodynamic limit. To get additional information on this mid gap state, we have plotted in Fig. 7 (d) the local magnetization in the sector $S^z_{\rm tot} = 1$. The local magnetization does not seem to decay exponentially. So it seems that the mid-gap state is not a well localized edge state.

Finally, the situation changes again for weakly coupled chains, in which case open boundary conditions with diagonal edges give rive to several mid-gap states (see Fig. 7 (c)). This can be understood as follows: In the limit of completely decoupled chains, there is an effective spin 1/2 at each end, i.e. 4 in total. So we expect to find 16 low-lying states in total. However, as soon as a small coupling between the chains is introduced, the two free spins $S = 1/2$ at one end of the ladder are expected to couple, leading to a finite excitation gap. This is confirmed by the magnetization, which is no longer localized at the edges of the chain (see Fig. 7 (d)) presumably because the spin-1/2 edges of the decoupled chains are bound into singlets.

CORRELATION LENGTH FROM LOCAL MAGNETIZATION

For the spin-1/2 ladder, when the system is in the strong rung coupling phase with one valence-bond singlet per rung, the correlation length can be obtained by fitting the decay of the lowest excitation energy with the size (see main text), or by fitting the decay of the local magnetization, as discussed in detail in this section. The local magnetization is defined as the expectation value of S^z_i in the ground state of the sector $S^z_{\rm tot} = 1$. The distribution of local magnetization is symmetric with respect to the vertical cut in the middle. Therefore only half the ladder can be taken into account for the fit. Moreover, in order to minimize the effect of the overlap of the left and right sides of the ladder, we do not consider the two rungs in the middle. Finally, we also did not include the magnetization of the first site to avoid the boundary effect. Then a log-linear plot of the local magnetization leads to a linear curve with a very good accuracy, from which a correlation length can easily be extracted (See Fig. 8).



Figure 7: (Color online) Low-energy spectra of the spin $S = 1$ ladder for different boundary conditions. In (a), we are in the rung singlet phase and we clearly see an edge states for OBC, diagonal edge. The fits use exponentially decaying functions in N. In (b) we are in the intermediate phase. (c) corresponds to two weakly coupled spin-1 chains. Valence-bond singlet sketches are attached to spectra (a) and (c). Dots denote spin-1/2, bold lines denote singlets, a single arrow indicates an isolated spin-1/2, and a double arrow stand for a spin-1. (d) Local magnetization in the sector $S^z_{\rm tot} = 1$. Squares and circles distinguish spins from top and bottom chains.



Figure 8: (Color online) Log-linear plot of the local magnetization for the 30-site spin-1/2 ladder with open boundary conditions and diagonal edges calculated in the ground state of the sector $S^z_{tot} = 1$. The correlation length is extracted from a linear fit.

PHASE TRANSITION AT $\theta = \pi$ AND $3\pi/2$

In the phase diagrams of the main text, we have indicated that the ferromagnetic phase extends from $\theta = \pi$ an $\theta = 3\pi/2$. That the ground state is ferromagnetic in this range is clear since both couplings are ferromagnetic. To prove that the ferromagnetic phase does not extend beyond this range, we have compared the ground state energies in all sectors of S^z_{tot}, which are expected to be completely degenerate only if the ground state is ferromagnetic (see Fig. 9).



Figure 9: (Color online) Ground state energies of a 16-site spin-1/2 ladder in the different sectors of S^z_{tot}. The ground state is completely degenerate, hence ferromagnetic, for $\theta \in (\pi, 3\pi/2)$.

[1] T. Hirano, H. Katsura, Y. Hatsugai, Phys.Rev. B **77**, 094431 (2008)